Table S1. The effects of treatment and time on photosynthetic gas exchange, chlorophyll fluorescence, photosynthetic pigments, and carbohydrates.

Two-way ANOVA was calculated using the formula: Variable ~ Treatment * Time + (1|Trial) + (1|Indiv) Treatment and time were included as categorical fixed factors; experimental replicate (Trial) and individual seedling (Indiv) were included as random factors. *P*-values in bold indicate statistical significance ($\alpha = 0.05$).

		Treatment		<u>Time</u>		$\underline{\mathbf{Treat} \times \mathbf{Time}}$	
Category	Parameter	${m F}$	P	${m F}$	P	${\it F}$	P
Photosynthetic	Anet	84.040	< 0.001	4.260	< 0.001	2.950	< 0.001
gas exchange	g_s	9.859	< 0.001	0.478	0.824	1.244	0.226
	$R_{\rm d}$	5.083	0.007	9.301	< 0.001	1.488	0.132
	Е	42.610	< 0.001	1.239	0.288	2.559	0.004
	IWUE	46.904	< 0.001	0.858	0.527	1.563	0.105
Chlorophyll	$F_{v}\!/F_{m}$	141.680	< 0.001	18.854	< 0.001	11.304	< 0.001
fluorescence	$\Phi_{ ext{PSII}}$	98.580	< 0.001	2.862	0.011	10.637	< 0.001
	NPQ	12.849	< 0.001	5.952	< 0.001	2.292	0.009
	NPQs	144.522	< 0.001	14.471	< 0.001	8.739	< 0.001
	1-qP	71.844	< 0.001	4.549	< 0.001	7.079	< 0.001
Photosynthetic	Total Chl	9.795	< 0.001	11.504	< 0.001	3.160	< 0.001
pigments	Car/Chl	12.940	< 0.001	4.758	< 0.001	1.467	0.139
	Chl a/b	7.917	< 0.001	0.941	0.467	3.200	< 0.001
	V+A+Z/Chl	34.125	< 0.001	4.127	< 0.001	2.314	0.009
	DEPS	61.462	< 0.001	21.187	< 0.001	11.718	< 0.001
Carbohydrates	Starch	9.397	< 0.001	46.018	< 0.001	3.788	< 0.001
	Soluble	2.916	0.057	23.846	< 0.001	1.139	0.333
	Sucrose	3.867	0.023	9.767	< 0.001	1.422	0.158
	Hexose	39.158	< 0.001	3.619	0.002	5.176	< 0.001
	Pinitol	2.116	0.123	14.114	< 0.001	1.533	0.115
	Raffinose	9.033	<0.001	5.417	<0.001	3.958	<0.001





Figure S1. Changes in total leaf protein content A) under controlled conditions in growth chambers in response to low temperature/ambient CO_2 (LTAC), elevated temperature/ambient CO_2 (ETAC), and elevated temperature/elevated CO_2 (ETEC) treatments; and B) over autumn under field conditions. Grey background indicates long photoperiod controls; white background indicates short photoperiod treatments. Bars represent the average of A) $n = 8-10 \pm SE$ biological replicates.



Quality

Figure S2. A) Predicted sequence of K-segment of the 16-kD dehydrin and B) alignment of peptides used to generate consensus. Consensus K-segment is outlined in red. Accession numbers for homologous proteins matched to each peptide are shown. Amino acids are color coded according to their chemical properties using the Lesk color scheme.




Figure S3. Response of shoot freezing tolerance to A) CO_2 level in seedlings grown under long photoperiod; B) low temperature/ambient CO_2 (LTAC), elevated temperature/ambient CO_2 (ETAC), and elevated temperature/elevated CO_2 (ETEC) in seedlings grown under short photoperiod. Grey background indicates long photoperiod controls; white background indicates short photoperiod treatments after 36 days. Grey vertical lines indicate midpoints representing LT_{50} . Letters, where present, indicate statistically different groups as determined by extra sum-of-squares F test.